

# Electrochemistry

## Electrochemical cells:

- Two types – Galvanic cell or voltaic cell, and Electrolytic cell

### Galvanic cell:

Converts the chemical energy of a spontaneous redox reaction into electrical energy

Daniell cell -

$$\Delta_r G^\ominus = -RT \ln K$$

The standard potential of a cell is given by

$$\Delta_r G^\ominus = -nFE_{\text{cell}}^\ominus$$

The potential of an individual half cell cannot be measured.

Relation between the standard potential of a cell and standard Gibbs energy:

$$\Delta_r G^\ominus = -nFE_{\text{cell}}^\ominus$$

Relation between standard Gibbs energy and the equilibrium constant:

$$\Delta_r G^\ominus = -RT \ln K$$

- Nernst equation:
  - Gives the concentration dependence of the potentials of the electrodes and the cells


For the electrode reaction

$$aA + bB \xrightarrow{n\text{e}^-} cC + dD$$

Nernst equation is given by

$$E_{\text{cell}} = E_{\text{cell}}^\ominus - \frac{RT}{nF} \ln \frac{[C]^c [D]^d}{[A]^a [B]^b}$$


Nernst equation is given by



Nernst equation is given by

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{RT}{nF} \ln \frac{[C]^c [D]^d}{[A]^a [B]^b}$$

- For a general electrochemical reaction of the type



Nernst equation is given by

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{RT}{nF} \ln \frac{[C]^c [D]^d}{[A]^a [B]^b}$$

### Conductance of electrolytic solutions:

Resistance,  $R = \rho \frac{l}{A}$

Where,

$l \rightarrow$  Length

$A \rightarrow$  Area of cross-section

$\rho \rightarrow$  Resistivity or specific resistance

Conductance,  $G = \frac{1}{R} = \frac{A}{\rho l} = \kappa \frac{A}{l}$

Where,  $\kappa$  Conductivity or specific conductance

The SI unit of conductance is  $\Omega^{-1}$  (siemens or mho).

The conductivity of an electrolyte depends upon

- nature of the solvent
- nature of the electrolyte added
- concentration of the electrolyte
- temperature

Molar conductivity,  $\Lambda_m = \frac{\kappa}{C}$

### Variation of conductivity

- For both strong and weak electrolytes, conductivity decreases with decrease in concentration.

### Variation of molar conductivity

- For both strong and weak electrolytes, molar conductivity increases with decrease in concentration.

**Limiting molar conductivity** – molar conductivity when concentration approaches zero

$$\text{Degree of dissociation, } \alpha = \frac{\Lambda_m}{\Lambda_m^0}$$

- Kohlrausch law of independent migration of ions:**

According to this law, for an electrolyte, the molar conductivity at infinite dilution is the sum of the contribution of the molar conductivity of the ions in which it dissociates.

### Electrolytic cells and electrolysis

$$1F = 96487 \text{ C mol}^{-1}$$

- Faraday's first law of electrolysis:** The amount of chemical reaction occurring at any electrode during the process of electrolysis by a current is proportional to the quantity of electricity passed through the electrolyte.
- Second law of electrolysis:** The amounts of different substances liberated when same quantity of electricity is passed through the electrolytic solution are proportional to their chemical equivalent weights.

Battery is a galvanic cell in which chemical energy of the redox reaction is converted into electrical energy.

Mainly two types:

- Primary batteries
- Secondary batteries
- Primary Batteries**

In primary batteries, reaction occurs only once.

After use over a period of time, these become dead and cannot be reused.

Examples: Dry cell (or Leclanche cell), Mercury cell

### Secondary Batteries

- Secondary batteries can be recharged again by passing current through them in the opposite direction.


- Examples: Lead storage battery, Nickel-cadmium cell

**Corrosion:** Oxidation of a metal by loss of electrons to oxygen and formation of oxides

### Corrosion of Iron

Known as rusting

The spot where oxidation takes place behaves as anode.



### Prevention of Corrosion

- Preventing the surface of the metal from coming in contact with atmosphere
- By covering the surface with paint or chemicals such as bis-phenol
- By covering the surface with other metals such as Sn, Zn, Mg.

**The Hydrogen Economy:** Based on electrochemical principles