

# Chapter 1 - Relations and Functions

## Definitions:

Let A and B be two non-empty sets, then a function f from set A to set B is a rule which associates each element of A to a unique element of B.

### ○ Relation

If  $(a, b) \in R$ , we say that a is related to b under the relation R and we write as  $a R b$

### ○ Function

It is represented as  $f: A \rightarrow B$  and function is also called mapping.

### ○ Real Function

$f: A \rightarrow B$  is called a real function, if A and B are subsets of R.

### ○ Domain and Codomain of a Real Function

Domain and codomain of a function f is a set of all real numbers x for which  $f(x)$  is a real number. Here, set A is domain and set B is codomain.

### ○ Range of a real function

f is a set of values  $f(x)$  which it attains on the points of its domain

## Types of Relations

- A relation R in a set A is called **Empty relation**, if no element of A is related to any element of A, i.e.,  $R = \emptyset \subset A \times A$ .
- A relation R in a set A is called **Universal relation**, if each element of A is related to every element of A, i.e.,  $R = A \times A$ .
- Both the empty relation and the universal relation are sometimes called **Trivial Relations**
- A relation R in a set A is called
  - **Reflexive**
    - if  $(a, a) \in R$ , for every  $a \in A$ ,
  - **Symmetric**
    - If  $(a_1, a_2) \in R$  implies that  $(a_2, a_1) \in R$ , for all  $a_1, a_2 \in A$ .
  - **Transitive**
    - If  $(a_1, a_2) \in R$  and  $(a_2, a_3) \in R$  implies that  $(a_1, a_3) \in R$ , for all  $a_1, a_2, a_3 \in A$ .
- A relation R in a set A is said to be an **equivalence relation** if R is reflexive, symmetric and transitive
- The set E of all even integers and the set O of all odd integers are subsets of Z satisfying following conditions:
  - All elements of E are related to each other and all elements of O are related to each other.
  - No element of E is related to any element of O and vice-versa.
  - E and O are disjoint and  $Z = E \cup O$ .
  - The subset E is called the equivalence class containing zero, Denoted by  $[0]$ .
  - O is the equivalence class containing 1 and is denoted by  $[1]$ .

- **Note**
  - $[0] \neq [1]$
  - $[0] = [2r]$
  - $[1] = [2r + 1], r \in \mathbb{Z}$ .
- Given an arbitrary equivalence relation  $R$  in an arbitrary set  $X$ ,  $R$  divides  $X$  into mutually disjoint subsets  $A_i$  called partitions or subdivisions of  $X$  satisfying:
  - All elements of  $A_i$  are related to each other, for all  $i$ .
  - No element of  $A_i$  is related to any element of  $A_j$ ,  $i \neq j$ .
  - $\bigcup A_i = X$  and  $A_i \cap A_j = \emptyset, i \neq j$ .
- The subsets  $A_i$  are called equivalence classes.

**Note:**

- Two ways of representing a relation
  - Roaster method
  - Set builder method
- If  $(a, b) \in R$ , we say that  $a$  is related to  $b$  and we denote it as  $a R b$ .


### Types of Functions

Consider the functions  $f_1, f_2, f_3$  and  $f_4$  given


- A function  $f: X \rightarrow Y$  is defined to be **one-one (or injective)**, if the images of distinct elements of  $X$  under  $f$  are distinct, i.e., for every  $x_1, x_2 \in X$ ,  $f(x_1) = f(x_2)$  implies  $x_1 = x_2$ . Otherwise,  $f$  is called **many-one**.

Example

- One- One Function





- Many-One Function




- A function  $f: X \rightarrow Y$  is said to be **onto (or surjective)**, if every element of  $Y$  is the image of some element of  $X$  under  $f$ , i.e., for every  $y \in Y$ , there exists an element  $x$  in  $X$  such that  $f(x) = y$ .

- $f: X \rightarrow Y$  is onto if and only if Range of  $f = Y$ .
- Eg:




- A function  $f: X \rightarrow Y$  is said to be **one-one and onto (or bijective)**, if  $f$  is both one-one and onto.
- Eg:



## Composition of Functions and Invertible Function

### Composite Function

- Let  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be two functions.
- Then the composition of  $f$  and  $g$ , denoted by  $g \circ f$ , is defined as the function  $g \circ f: A \rightarrow C$  given by  $g \circ f(x) = g(f(x)), \forall x \in A$ .




- Eg:
  - Let  $f: \{2, 3, 4, 5\} \rightarrow \{3, 4, 5, 9\}$  and  $g: \{3, 4, 5, 9\} \rightarrow \{7, 11, 15\}$  be functions
  - Defined as  $f(2) = 3, f(3) = 4, f(4) = f(5) = 5$  and  $g(3) = g(4) = 7$  and  $g(5) = g(9) = 11$ .
  - Find  $g \circ f$ .
  - Solution
    - $g \circ f(2) = g(f(2)) = g(3) = 7$ ,
    - $g \circ f(3) = g(f(3)) = g(4) = 7$ ,
    - $g \circ f(4) = g(f(4)) = g(5) = 11$  and
    - $g \circ f(5) = g(5) = 11$
- It can be verified in general that  $g \circ f$  is one-one implies that  $f$  is one-one. Similarly,  $g \circ f$  is onto implies that  $g$  is onto.

- While composing  $f$  and  $g$ , to get  $gof$ , first  $f$  and then  $g$  was applied, while in the reverse process of the composite  $gof$ , first the reverse process of  $g$  is applied and then the reverse process of  $f$ .
- If  $f: X \rightarrow Y$  is a function such that there exists a function  $g: Y \rightarrow X$  such that  $gof = IX$  and  $fog = IY$ , then  $f$  must be one-one and onto.

### Invertible Function

- A function  $f: X \rightarrow Y$  is defined to be invertible, if there exists a function  $g: Y \rightarrow X$  such that  $gof = IX$  and  $fog = IY$ . The function  $g$  is called the inverse of  $f$
- Denoted by  $f^{-1}$ .



- Thus, if  $f$  is invertible, then  $f$  must be one-one and onto and conversely, if  $f$  is one-one and onto, then  $f$  must be invertible.

### Theorem 1

- If  $f: X \rightarrow Y$ ,  $g: Y \rightarrow Z$  and  $h: Z \rightarrow S$  are functions, then
  - $h \circ (g \circ f) = (h \circ g) \circ f$ .
- Proof

We have

- $h \circ (g \circ f) (x) = h(g \circ f (x)) = h(g(f(x))), \forall x \in X$
- $(h \circ g) \circ f (x) = h \circ g (f(x)) = h (g (f(x))), \forall x \in X$ .

Hence,  $h \circ (g \circ f) = (h \circ g) \circ f$

### Theorem 2

- Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be two invertible functions.
  - Then  $gof$  is also invertible with  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$
- Proof
  - To show that  $gof$  is invertible with  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ , it is enough to show that  $(f^{-1} \circ g^{-1}) \circ (g \circ f) = IX$  and  $(g \circ f) \circ (f^{-1} \circ g^{-1}) = IZ$ .  
 Now,  $(f^{-1} \circ g^{-1}) \circ (g \circ f) = ((f^{-1} \circ g^{-1}) \circ g) \circ f$ , by Theorem 1  
 $= (f^{-1} \circ (g^{-1} \circ g)) \circ f$ , by Theorem 1  
 $= (f^{-1} \circ IX) \circ f$ , by definition of  $g^{-1}$   
 $= IX$

Similarly, it can be shown that  $(g \circ f) \circ (f^{-1} \circ g^{-1}) = IZ$

### Binary Operations

#### Definitions:

- A binary operation  $*$  on a set  $A$  is a function  $*: A \times A \rightarrow A$ . We denote  $*(a, b)$  by  $a * b$ .

- A binary operation  $*$  on the set  $X$  is called commutative, if  $a * b = b * a$ , for every  $a, b \in X$ .
- A binary operation  $* : A \times A \rightarrow A$  is said to be associative if  $(a * b) * c = a * (b * c)$ ,  $\forall a, b, c \in A$ .
- A binary operation  $* : A \times A \rightarrow A$ , an element  $e \in A$ , if it exists, is called identity for the operation  $*$ , if  $a * e = a = e * a$ ,  $\forall a \in A$ .
  - Zero is identity for the addition operation on  $R$  but it is not identity for the addition operation on  $N$ , as  $0 \notin N$ .
  - Addition operation on  $N$  does not have any identity.
  - For the addition operation  $+ : R \times R \rightarrow R$ , given any  $a \in R$ , there exists  $-a$  in  $R$  such that  $a + (-a) = 0$  (identity for '+')  $= (-a) + a$ .
  - For the multiplication operation on  $R$ , given any  $a \neq 0$  in  $R$ , we can choose  $\frac{1}{a}$  such that  $a \times \frac{1}{a} = 1$  (identity for '×')  $= 1 = \frac{1}{a} \times a$
- A binary operation  $* : A \times A \rightarrow A$  with the identity element  $e$  in  $A$ , an element  $a \in A$  is said to be invertible with respect to the operation  $*$ , if there exists an element  $b$  in  $A$  such that  $a * b = e = b * a$  and  $b$  is called the inverse of  $a$  and is denoted by  $a^{-1}$